arXiv:1401.4511v1 [astro-ph.HE] 18 Jan 2014

Fermi LAT detection of two high Galactic latitude gamma-ray sources, Fermi J1049.7+0435 and J1103.2+1145

Masaki Nishimichi, Takeshi Okuda and Masaki Mori

Department of Physical Sciences, Ritsumeikan University, Kusatsu, Shiga 525-8577, Japan

Philip G. Edwards

CSIRO Astronomy and Space Science, PO Box 76, Epping NSW 1710, Australia

Jamie Stevens

CSIRO Astronomy and Space Science, Locked Bag 194, Narrabri NSW 2390, Australia

Received _____; accepted _____

To appear in Astrophysical Journal

ABSTRACT

During a search for gamma-ray emission from NGC 3628 (Arp 317), two new unidentified gamma-ray sources, Fermi J1049.7+0435 and J1103.2+1145 have been discovered (Nishimichi & Mori 2013). The detections are made in data from the Large Area Telescope (LAT), on board the Fermi Gamma-ray Space Telescope, in the 100 MeV to 300 GeV band during the period between 2008 August 5 and 2012 October 27. Neither is coincident with any source listed in the 2FGL catalogue (Nolan et al. 2012). Fermi J1049.7+0435 is at Galactic coordinates $(l, b) = (245.34°, 53.27°)$, $(\alpha_{J2000}, \delta_{J2000}) = (162.43°, 4.60°)$. Fermi J1103.2+1145 is at Galactic coordinates $(l, b) = (238.85°, 60.33°)$, $(\alpha_{J2000}, \delta_{J2000}) = (165.81°, 11.75°)$. Possible radio counterparts are found for both sources, which show flat radio spectra similar to other Fermi LAT detected AGN, and their identifications are discussed.

Subject headings: BL Lacertae objects: general — galaxies: active — gamma-rays: galaxies

1. Introduction

The Second Fermi LAT source catalog (Nolan et al. 2012) includes as many as 1,873 sources, but initial attempts to identify counterparts at other wavelengths resulted in 575 sources remaining unidentified. The 2FGL catalog is based on the first 24 months of LAT observation since its launch in 2008, but the LAT has now accumulated more than 5 years of high-energy gamma-ray data almost flawlessly, presenting the possibility of finding new sources which were too faint to be detected in the first two years of data or showed flaring activity after the catalog was created.

In this paper we report on two new gamma-ray sources serendipitously discovered in the constellation Leo and discuss possible counterparts based on radio observations.

2. Analysis

Our original aim was to search for gamma-ray emission from NGC 3628 (Arp 317), one of the three galaxies called the 'Leo Triplet', for which possible starburst activity has been reported based on XMM observations (Tüllman et al. 2006). Five years of archival data of Fermi LAT has been analyzed using the Fermi Science Tools supplied by Fermi Science Support Center (FSSC (2013), Fermi Science Tools v9r23p1). The energy range used in the present analysis was from 100 MeV to 300 GeV. 'Source' class events detected at zenith angles smaller than 100° were used for analysis, assuming 'P7SOURCE_V6' instrument response function along with the standard analysis pipeline suggested by FSSC. The significance of gamma-ray signal has been estimated by maximum likelihood method with a help of the `gtlike` program (which we used in the binned mode) included in the tools. The data periods for this studies span from 2008 August 4 to 2012 October 27.

For NGC 3628 (Arp 317), the test statistic, TS, returned by `gtlike` is consistent with

zero, indicating there is no evidence of gamma-ray emission. Thus we calculated upper limits to gamma-ray flux from NGC 3628 of $1.4\,(1.3) \times 10^{-9}$ cm^{-2}s^{-1}, at 95% C.L., above 100 MeV for the period 2008 August 4 to 2010 July 31 (2010 July 31 to 2012 October 27). This is translated to gamma-ray luminosity upper limit of $2.5\,(2.3) \times 10^{43}$ erg s^{-1} assuming the distance of 12 Mpc which is derived as the median of 8 measurements, which range from 6.7 to 14.2 Mpc (NED 2012).

During the study of NGC 3628, we noticed two rather bright gamma-ray sources in the field of view centered on NGC 3628 and within a radius of 15° (Nishimichi & Mori 2013). They are not coincident with any source listed in the 2FGL catalogue (Nolan et al. 2012) nor in the 3EG catalogue (Hartman et al. 1999). Figure 1 shows a gamma-ray countmap of this area. The positions for these sources were estimated using the `gttsmap` program which calculates the TS value assuming an unknown source at various positions in the field-of-view of interest, and the maximum TS values were obtained for positions shown in the Table 1. The errors of the positions are conservatively estimated as the radius at which the TS value drops to the half value. Figures 2 and 3 show the TS maps around the new sources for the half-year period where the TS value takes the maximum, which are consistent with the assumption of point sources. The average fluxes, the power-law indices, and the TS value for the whole analyzed period (fron 2008 August 27 to October 10, 2013) are: $(3.68 \pm 0.36) \times 10^{-8}$ cm^{-2}s^{-1} (> 100 MeV), -2.60 ± 0.07 and $TS = 332$ for J1049.7+0435; $(2.35 \pm 0.30) \times 10^{-8}$ cm^{-2}s^{-1} (> 100 MeV), -2.55 ± 0.09 and $TS = 196$ for J1103.2+1145.

Figures 4 and 5 show the time variation of gamma-ray fluxes of the newly detected sources in half-year bins. For these plots we added data until 2013 October 10. One can see in the first two years their fluxes are below the detection threshold ($TS < 25$), which is why they are not listed in the 2FGL catalog based on data over the similar period (Nolan et al.



Fig. 1.— Gamma-ray countmap around the NGC 3628 region. The map is created in 0.1° grid and smoothed for the data during 2008 August 05 to 2013 July 03. 2FGL sources are annotated, and two new gamma-ray sources are marked as 'Unknown_1' (J1103.2+1145) and 'Unknown_2' (J1049.7+0435).



Fig. 2.— Test-statistics map around the J1049.7+0435 region for the period from 2011 October 27 to 2012 April 27, where the TS value is the highest ($TS = 222$).



Fig. 3.— Test-statistics map around the J1103.2+1145 region for the period from 2013 April 27 to 2013 October 27, where the TS value is the highest ($TS = 197$).

Table 1: Best positions of new sources

Name	α_{J2000} (deg)	δ_{J2000} (deg)	$\ell^{\rm II}$ (deg)	$b^{\rm II}$ (deg)	error radius (arcmin)
J1049.7+0435	162.43	4.60	245.34	53.27	51
J1103.2+1145	165.81	11.75	238.85	60.33	66

2012). The first of our two sources has subsequently been detected in the Fermi All-sky Variability Analysis (Ackermann et al. 2013) and catalogued as 1FAV J1051+04.

3. Discussion

The variability of both sources display at gamma-ray energies suggests they are more likely to be AGN than members of other populations of identified gamma-ray sources (Nolan et al. 2003). The gamma-ray spectral indices are more consistent with those of flat spectrum radio quasars (FSRQs) than of BL Lac objects (Ackermann et al. 2011): FSRQs are on average found to be more variable than BL Lac objects (Nolan et al. 2003; Ackermann et al. 2011).

Mattox et al. (1997, 2001) showed that (extragalactic) gamma-ray sources were more likely to be associated with brighter radio sources, and in particular those sufficiently compact to be detectable in VLBI observations. Radio compactness is generally associated with a flatter radio spectrum, resulting from synchtrotron self-absorption at lower frequencies, which is also a characteristic of Fermi-detected AGN (Abdo et al. 2010). (Although, as noted by, e.g., Edwards (2005), radio spectral indices determined from single dish observations are affected by steeper-spectrum radio lobes in some sources, which disguise the presence of a flat-spectrum radio core.)

We have, therefore, searched for potential counterparts in the Green Bank 6-cm (GB6, Gregory et al. (1996)) catalog and determined spectral indices between 20 cm and 6cm using the NRAO VLA Sky Survey (NVSS, Condon et al. (1998)) catalog.

The closest GB6 radio source to J1049.7+0435 is GB6 J1050+0432, with an angular separation of 7 arcmin. The source has a flux density of 99±10 mJy at 4.8 GHz, and the corresponding 20 cm source, NVSS J105010+043251, has a flux density of 101.2 mJy,



Fig. 4.— Time variation of gamma-ray flux of J1049.7+0435 in half-year bins for the period from 2008 August 5 to 2013 October 10. Triangles are upper limits (95% C.L.).



Fig. 5.— Time variation of gamma-ray flux of J1103.2+1145 in half-year bins for the period from 2008 August 5 to 2013 October 10. Triangles are upper limits (95% C.L.).

yielding a spectral index α (where $S \propto \nu^{+\alpha}$) of 0.0. Two fractionally brighter GB6 sources have both larger angular offsets and signficantly **steeper** spectra: GB6 J1049+0505, 113 mJy, 30 arcmin separation, α=−0.8; GB6 J1051+0449, 101 mJy, 29 arcmin separation, α=−0.9. We note that the GB6 and NVSS flux densities were made some years apart, and so these spectral indices should be taken as representative values rather than absolute measurements. As this declination range is also covered by the Parkes-MIT-NRAO equatorial survey (Griffith et al. 1995), we can compare the GB6 value with that PMN J1050+0432, which has a 4.8 GHz flux density of 98±12 mJy.

For J1105.2+1145, the closest GB6 source is GB6 J1103+1158, with an angular separation of 14 arcmin. The source has a 4.8 GHz flux density of 306±27 mJy, with the corresponding 20cm source, NVSS J110303+115816, having a flux density of 262.6 mJy, resulting in a spectral index of 0.1. Other relatively bright GB6 sources in the area are further away and with steeper spectral indices: GB6 J1103+1114, 116 mJy, 31 arcmin, α=−0.7; GB6 J1104+1103, 277 mJy, 46 arcmin, α=−0.8. A Seyfert 1 galaxy, Mrk 728, is 0.89 deg from J1103.2+1145 and is not likely the counterpart. GB6 J1103+1158 corresponds to the quasar SDSS 110303.52+115816.5, which lies at a redshift of 0.912 (Schneider et al. 2007). Furthermore, the quasar has been detected in the VLBA Calibrator Survey VLBI observations (Petrov et al. 2005), confirming the presence of a compact core in this radio-loud AGN.

Catalogued radio positions and flux densities for the two sources are tabulated in Table 2.

We have additionally made snap-shot observations of J1049.7+0435 and J1103.2+1145 (at their NVSS positions) with the Australia Telescope Compact Array at several epochs, as part of an on-going program to monitor gamma-ray sources (Stevens et al. 2012) with the measured flux densities are listed in Table 3. The observations at 17 GHz and 38 GHz were

Table 2: Possible radio counterparts,

Gamma-ray source	catalog (frequency)	RA (J2000)	Dec (J2000)	Radio flux density (mJy)
J1049.7+0435	NVSS (1.4 GHz)	10 50 10.06	+04 32 51.3	101.2
	GB6 (4.8 GHz)	10 50 08.6	+04 32 37	99
J1103.2+1145	NVSS (1.4 GHz)	11 03 03.55	+11 58 16.6	262
	GB6 (4.8 GHz)	11 03 03.7	+11 58 20	306

preceded by a pointing scan on a nearby bright compact source to refine the global pointing model. Data were processed in Miriad in the standard manner. Flux density calibration was bootstrapped to the standard ATCA flux density calibrator, PKS 1934−638. Errors are conservatively estimated as 5% at lower frequencies and 10% at highest frequencies, where these include statistical and systematic errors, with the latter dominating.

GB6 J1050+0432 has brightened considerably, by a factor of 2.7, since the GB6 and PMN observations (which date back to the late 1980s and early 1990s), and has an inverted spectrum with $\alpha \sim 0.25$, strengthening the case for an association with J1049.7+0435. Note also the increased gamma-ray flux in the latest half year (Fig.4).

GB6 J1103+1158 is a little fainter than the catalogued GB6 value, however the ATCA observations confirm that the spectral index remains flat, at $\alpha \sim -0.1$, up to 38 GHz. There is no evidence of significant variability over the 4 months spanned by these observations, however comparison with the GB6 flux density indicates the presence of longer timescale variability.

We have also examined the ASDC Sky Explorer (ASDC, ASDC (2012)) and NASA/IPAC Extragalactic Database (NED, NED (2012)) for other possible counterparts,

Table 3: ATCA radio observations (unit: mJy). See text for details.

Gamma-ray source	Epoch	Array	5.5 GHz	9.0 GHz	17 GHz	38 GHz
GB6 J1050+0432	2013 Oct 20	H214	276	311	371	
GB6 J1103+1158	2013 May 10	6C	254	238	216	230
	2013 Aug 20	H168	254	245		
	2013 Sep 8	1.5A	246	230		

but we did not find any good candidates nearer than radio sources mentioned above.

In the light of the above facts, we tentatively identify both gamma-ray sources with the radio sources mentioned above. Petrov et al. (2013) make a detailed consideration of the utility of radio observations in finding counterparts to unidentified Fermi sources. The associations proposed here would be strengthened by improved gamma-ray localisations, and/or evidence of contemporaneous multi-wavelength flaring, and, in the case of GB6 J1050+0432, with VLBI observations to determine whether the source contains a compact, parsec-scale, radio core.

4. Conclusions

A search for gamma-rays from NGC 3628 (Arp 317), for which possible starburst activity has been reported, found no evidence for >100 MeV emission. However, two new GeV sources, Fermi J1049.7+0435 and J1103.2+1145, have been found near the Leo Triplet region using Fermi-LAT archival data spanning 5 years. The fluxes for both sources increase over the 5 yr period: thus they are not included in 2FGL catalog. Their flux variability and spectral indices are compatible with those of gamma-ray detected AGN. Based on angular separation, radio flux density and spectral index, we associate J1049.7+0435

with GB6 J1050+0432, and J1103.2+1145 with the quasar GB6 J1103+1158. Further multiwavelength studies are required to confirm these identifications.

Finally, looking forward to the release of the third Fermi LAT catalog, we note that the 15 deg radius (0.2 sr) field studied here has yielded two previously uncataloged sources. Although the small number statistics result in extrapolations having large uncertainties, this suggests (ignoring further improvements in analysis software and background models) there might be $\sim$100 new extra-galactic sources with $|b| > 10^\circ$ in the 3FGL catalog.

This work is supported in part by the Grant-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan, No. 22540315. The Australia Telescope Compact Array is part of the Australia Telescope National Facility which is funded by the Commonwealth of Australia for operation as a National Facility managed by CSIRO. Leonid Petrov is acknowledged for his independent suggestion of GB6 J1103+1158 as the possible counterpart to J1103.2+1145.

REFERENCES

Abdo, A.A., Ackermann, M., Ajello, M. et al., ApJ, 2010, 715, 429

Ackermann, M., Ajello, M., Allafort, A., et al. 2011, ApJ, 743, 171

Ackermann, M., Ajello, M., Allafort, A., et al. 2012, ApJ, 753, 83

Ackermann, M., Ajello, M., Albert, A., et al. 2013, ApJ, 771, 57

ASI Science Data Center (ASDC), http://www.asdc.asi.it/

Condon, J. J., Cotton, W. D., Greisen, E. W., et al. 1998, AJ, 115, 1693

Edwards, P. G. 2005, Ap&SS, 297, 31

Fermi Science Support Center (FSSC), http://fermi.gsfc.nasa.gov/ssc/

Gregory, P. C., Scott, W. K., Douglas, K., & Condon, J. J. 1996, ApJS, 103, 427

Griffith, M. R., Wright, A. E., Burke, B. F., & Ekers, R. D. 1995, ApJS, 97, 347

Hartman, R. C., Bertsch, D. L., Bloom, S. D., et al. 1999, ApJS, 123, 79

Mattox, J. R., Schachter, J., Molnar, L., Hartman, R. C., & Patnaik, A. R. 1997, ApJ, 481, 95

Mattox, J. R., Hartman, R. C., & Reimer, O. 2001, ApJS, 135, 155

NASA/IPAC Extragalactic Database (NED), http://ned.ipac.caltech.edu/

Nishimichi, M. & Mori, M., 2013, Astronomer's Telegram, #4949

Nolan, P. L., Tompkins, W. F., Grenier, I. A., & Michelson, P. F. 2003, ApJ, 597, 615

Nolan, P.L., Abdo, A.A., Ackermann, M. et al., ApJS, 2012, 199, 31

Petrov, L., Kovalev, Y. Y., Fomalont, E., & Gordon, D. 2005, AJ, 129, 1163

Petrov, L., Mahony, E. K., Edwards, P. G., et al. 2013, MNRAS, 432, 1294

Schneider, D. P., Hall, P. B., Richards, G. T., et al. 2007, AJ, 134, 102

Stevens, J., Edwards, P. G., Ojha, R., et al. 2012, Fermi & Jansky Proceedings — eConf C1111101, arXiv:1205.2403

Tüllman, R., Pietsch, W., Rossa, J. et al., A&A, 2006, 448, 43

This manuscript was prepared with the AAS LaTeX macros v5.2.